arXiv:1405.7147v1 [math.CO] 28 May 2014

NEW EXTREMAL BINARY SELF-DUAL CODES FROM $\mathbb{F}_4 + u\mathbb{F}_4$-LIFTS OF QUADRATIC CIRCULANT CODES OVER $\mathbb{F}_4$

ABIDIN KAYA, BAHATTIN YILDIZ, AND IRFAN SIAP

Abstract. In this work, quadratic double and quadratic bordered double circulant constructions are applied to $\mathbb{F}_4 + u\mathbb{F}_4$ as well as $\mathbb{F}_4$, as a result of which extremal binary self-dual codes of length 56 and 64 are obtained. The binary extension theorems as well as the ring extension version are used to obtain 7 extremal self-dual binary codes of length 58, 24 extremal self-dual binary codes of length 66 and 29 extremal self-dual binary codes of length 68, all with new weight enumerators, to update the list of all the known extremal self-dual codes in the literature.

1. Introduction

Double circulant, bordered double circulant and recently four circulant constructions are some of the well known methods by which self-dual codes especially the extremal ones are constructed and classified. While in the former two, the exact conditions when the resulting codes are self-dual are not known, in the four circulant case a necessary and sufficient condition is given for the resulting code to be self-dual. Another common theme related to binary self-dual codes has been using extension theorems to obtain self-dual codes of length $n+2$ from self-dual codes of length n.

Recently certain binary rings (rings of characteristic 2) have been successfully used to obtain many new extremal binary self-dual codes using the construction methods mentioned above and extension theorems. Some of the examples of these constructions can be found in [9, 10, 11, 12], etc.

Gaborit, in [6] combined the double and bordered double circulant constructions with quadratic residues modulo a prime to construct self-dual codes of certain lengths over finite fields. The construction was called quadratic double and quadratic bordered double circulant construction. In this case also, the conditions for the self duality of the constructed codes were expressed explicitly.

Gaborit's method over fields is extended to the rings by the authors in [14] for the ring $\mathbb{F}_2 + u\mathbb{F}_2 + u^2\mathbb{F}_2$ with $u^3 = u$ to obtain a considerable number of new extremal binary self-dual codes of length 68.

In a recent work ([15]), it was shown that the extension theorems for binary self-dual codes can also be extended to hold for self-dual codes over binary rings.

In this work, we combine all the above mentioned methods for some special rings, that are the rings $\mathbb{F}_{2^m} + u\mathbb{F}_{2^m}$ for $m = 1, 2$, where $u^2 = 0$. These rings have been

2000 *Mathematics Subject Classification.* Primary 94B05, 94B99; Secondary 11T71, 13M99.

Key words and phrases. quadratic residue codes, extremal self-dual codes, Gray maps, quadratic double-circulant codes.

studied in [1] and [16], where it was shown that they are endowed with a duality-preserving Gray map that allows us to construct binary self-dual codes from self-dual codes over $\mathbb{F}_{2^m}+u\mathbb{F}_{2^m}$. To elaborate on the main focus of the current work, we have combined the extension theorem for rings with quadratic double and quadratic bordered double circulant constructions over the rings $\mathbb{F}_2+u\mathbb{F}_2$ and $\mathbb{F}_4+u\mathbb{F}_4$ and using these methods, we have been able to construct a considerable number of new extremal binary self-dual codes of lengths 58, 66 and 68. The precise numbers of new codes obtained from the constructions that will be explained in subsequent chapters are as follows: 7 extremal binary self-dual codes of length 58 with new weight enumerators in $W_{58,2}$; 24 extremal binary self-dual codes of length 66 with new weight enumerators in $W_{66,3}$; and 29 extremal binary self-dual codes of length 68 with new weight enumerators in $W_{68,2}$.

The rest of the paper is organized as follows. In Section 2, we give the preliminaries about the rings $\mathbb{F}_2+u\mathbb{F}_2$, $\mathbb{F}_4+u\mathbb{F}_4$, binary self-dual codes and the extension theorems. Section 3 includes the quadratic double and bordered double circulant constructions for self-dual codes over $\mathbb{F}_4+u\mathbb{F}_4$ and $\mathbb{F}_4$ as well as the lifts of self-dual codes over $\mathbb{F}_4$. This leads to several extremal binary self-dual codes of lengths 56 and 64, which will form a basis for the extensions to be applied in Section 4. Extension theorems of various forms are applied in Section 4 to the extremal codes obtained in Section 3, as a result of which numerous new extremal binary self-dual codes of lengths 58, 66 and 68 are obtained. We conclude with remarks and possible directions for future research.

2. Preliminaries

Let $\mathbb{F}_4$ be the finite field of four elements, in other words $\mathbb{F}_4=\mathbb{F}_2(\omega)$ where ω is a root the unique irreducible binary quadratic polynomial x^2+x+1. The ring $\mathbb{F}_4+u\mathbb{F}_4$ defined via $u^2=0$ can be viewed as an extension of $\mathbb{F}_2, \mathbb{F}_4$ or $\mathbb{F}_2+u\mathbb{F}_2$. Self-dual codes over a ring of characteristic 2 are called Type II if the Lee weights are multiples of 4 and Type I otherwise. Type II codes over $\mathbb{F}_4+u\mathbb{F}_4$ is studied in [16], the results later generalized to the ring $\mathbb{F}_{2^m}+u\mathbb{F}_{2^m}$ [1]. For more details we refer to [16, 1]. Recently, four circulant codes over $\mathbb{F}_4+u\mathbb{F}_4$ have been studied in [15]. An $(\mathbb{F}_4+u\mathbb{F}_4)$-submodule C of $(\mathbb{F}_4+u\mathbb{F}_4)^n$ is called a linear code of length n over $\mathbb{F}_4+u\mathbb{F}_4$. Let $x=(x_1,x_2,\ldots,x_n)$ and $y=(y_1,y_2,\ldots,y_n)$, the dual $C^\perp$ of the code C is defined for the Euclidean inner product $\langle x,y\rangle=\sum x_iy_i$ as

$$C^\perp=\{x\in(\mathbb{F}_4+u\mathbb{F}_4)^n \mid \langle x,y\rangle=0 \text{ for all } y\in C\}.$$

C is said to be self-dual if $C=C^\perp$. Rains finalized the upper bound for the minimum distance d of a binary self-dual code of length n in [17] as $d\leq 4\lfloor\frac{n}{24}\rfloor+6$ if $n\equiv 22 \pmod{24}$ and $d\leq 4\lfloor\frac{n}{24}\rfloor+4$, otherwise. A self-dual binary code is called *extremal* if it meets the bound. Throughout the text we have used the following Gray maps from [16];

$$\begin{array}{l|l} \psi:(\mathbb{F}_4+u\mathbb{F}_4)^n\to(\mathbb{F}_2+u\mathbb{F}_2)^{2n} & \phi:(\mathbb{F}_2+u\mathbb{F}_2)^n\to\mathbb{F}_2^{2n} \\ a\omega+b\overline{\omega}\mapsto(a,b),\ \ a,b\in(\mathbb{F}_2+u\mathbb{F}_2)^n & a+bu\mapsto(b,a+b),\ \ a,b\in\mathbb{F}_2^n. \end{array}$$

The maps preserve orhogonality, therefore the Gray images of self-dual codes are self-dual.

Quadratic double circulant (QDC) codes are introduced in [6]. Let $Q_p(a,b,c)$ be a circulant matrix with first row r based on quadratic residues modulo p defined as $r[1]=a$, $r[i+1]=b$ if i is a quadratic residue and $r[i+1]=c$ if i is a quadratic

non-residue, we state the special case of the main theorem from [6] where p is an odd prime;

Theorem 2.1. ([6]) *Let p be an odd prime and let $Q_p(a,b,c)$ be a quadratic residue circulant matrix with a, b and c as the elements of the commutative ring R with identity. If $p = 4k+1$ then*

$$\begin{aligned} & Q_p(a,b,c)\, Q_p(a,b,c)^T \\ = \;& Q_p\left(a^2 + 2k\left(b^2+c^2\right), 2ab - b^2 + k(b+c)^2, 2ac - c^2 + k(b+c)^2\right). \end{aligned}$$

If $p = 4k+3$ then

$$\begin{aligned} & Q_p(a,b,c)\, Q_p(a,b,c)^T \\ = \;& Q_p(a^2 + (2k+1)\left(b^2+c^2\right), ab + ac + k\left(b^2+c^2\right) + (2k+1)bc, \\ & ab + ac + k\left(b^2+c^2\right) + (2k+1)bc). \end{aligned}$$

Definition 2.2. The code generated by $P_p(a,b,c) = \left[\; I_p \;\middle|\; Q_p(a,b,c) \;\right]$ over R is called a quadratic pure double circulant code and is denoted by $\mathcal{P}_p(a,b,c)$. In a similar way, the code generated by

$$B_p(a,b,c,\lambda,\beta,\gamma) = \left[\; I_{p+1} \;\middle|\; \begin{array}{cc} \lambda & \beta \times \mathbf{1} \\ \gamma \times \mathbf{1}^T & Q_p(a,b,c) \end{array} \;\right],$$

where $\mathbf{1}$ is the all 1 vector of length p, is called a bordered quadratic double circulant code and is denoted by $\mathcal{B}_p(a,b,c,\lambda,\beta,\gamma)$.

In order to define an $(\mathbb{F}_4 + u\mathbb{F}_4)$-lift of a code over $\mathbb{F}_4$ we need the projection $\mu : (\mathbb{F}_4 + u\mathbb{F}_4)^n \mapsto \mathbb{F}_4^n$ defined as $\mu\left(\overline{a} + \overline{b}u\right) = \overline{a}$.

Theorem 2.3. ([11]) ($\mathbb{F}_2$ *replaced by* $\mathbb{F}_4$) *Let C be a self-dual code of length n over $\mathbb{F}_4 + u\mathbb{F}_4$ then $\mu(C)$ is a self-orthogonal code of length n over $\mathbb{F}_4$.*

Definition 2.4. Let C and D be codes of length n over $\mathbb{F}_4 + u\mathbb{F}_4$ and $\mathbb{F}_4$, respectively. Then, C is said to be a lift of D if $\mu(C) = D$.

Throughout the text we consider quadratic double circulant codes over $\mathbb{F}_4$, their lifts to $\mathbb{F}_4 + u\mathbb{F}_4$, $\mathbb{F}_2 + u\mathbb{F}_2$ and $\mathbb{F}_2$ and ring extensions of the related self-dual codes. For extensions the following theorems from [15] are used. In the sequel, we let S to be a commutative ring of characteristic 2 with identity.

Theorem 2.5. ([15]) *Let C be a self-dual code over S of length n and $G = (r_i)$ be a $k \times n$ generator matrix for C, where r_i is the i-th row of G, $1 \le i \le k$. Let c be a unit in S such that $c^2 = 1$ and X be a vector in S^n with $\langle X, X\rangle = 1$. Let $y_i = \langle r_i, X\rangle$ for $1 \le i \le k$. Then the following matrix*

$$\left[\begin{array}{cc|c} 1 & 0 & X \\ \hline y_1 & cy_1 & r_1 \\ \vdots & \vdots & \vdots \\ y_k & cy_k & r_k \end{array}\right],$$

generates a self-dual code D over S of length $n+2$.

Another extension method which can be applied to generator matrices in standard form is as follows.:

Theorem 2.6. ([15]) *Let C be a self-dual code generated by $G = [I_n|A]$ over S. If the sum of the elements in i-th row of A is r_i then the matrix:*

$$G^* = \left[\begin{array}{cc|ccccc} 1 & 0 & x_1 & \dots & x_n & 1 & \dots & 1 \\ \hline y_1 & cy_1 & & & & & & \\ \vdots & \vdots & & & I_n & & A & \\ y_n & cy_n & & & & & & \end{array}\right],$$

where $y_i = x_i + r_i$, c is a unit with $c^2 = 1$, $X = (x_1, \dots, x_n)$ and $\langle X, X\rangle = 1 + n$, generates a self-dual code C^* over S.

3. Quadratic double circulant codes over $\mathbb{F}_4 + u\mathbb{F}_4$

We investigate QDC codes over $\mathbb{F}_4 + u\mathbb{F}_4$ and the $(\mathbb{F}_4 + u\mathbb{F}_4)$-lifts of QDC codes over $\mathbb{F}_4$. Note that a lift of a self-dual code may not be self-dual. But considering the $(\mathbb{F}_4 + u\mathbb{F}_4)$-lifts of a self-dual code over $\mathbb{F}_4$ reduces the workload remarkably. For instance, there are 8 free elements for a random double circulant code of length 16 over $\mathbb{F}_4 + u\mathbb{F}_4$ which results in $16^8 = 4294967296$ possibilities. On the other hand, if we consider the lifts of a unique double circulant code of length 16 over $\mathbb{F}_4$ the number of possibilities reduces to $4^8 = 65536$. This is because for a fixed element $a \in \mathbb{F}_4$ there are 4 possible lifts $a + bu \in \mathbb{F}_4 + u\mathbb{F}_4$ where b is an arbitrary element of $\mathbb{F}_4$. Results are obtained by computational algebra system MAGMA, for more details we refer the reader to [2]. By using these constructions we obtain some binary self-dual codes of lengths 56 and 64.

3.1. Quadratic double circulant codes over $\mathbb{F}_4+u\mathbb{F}_4$.

QDC codes over $\mathbb{F}_4+u\mathbb{F}_4$ is a large family of self-dual codes. In the following theorem we list some of these which correspond to self-dual codes.

Theorem 3.1. *Let p be a prime then the following holds:*
if $p = 8k + 7$ then the codes

$$\mathcal{P}_p\left(u, 1+\omega, \omega+u\omega\right) \text{ and } \mathcal{B}_p\left(1+u+u\omega, \omega+u\omega, 1+\omega, u, 1+u\omega, 1+u\omega\right)$$

are Type II codes over $\mathbb{F}_4 + u\mathbb{F}_4$,
if $p = 8k + 3$ then the codes

$$\mathcal{B}_p\left(1, u+u\omega, 1, u+u\omega, 1+u+u\omega, 1+u+u\omega\right) \text{ and}$$
$$\mathcal{B}_p\left(1+\omega, \omega+u\omega, 1+\omega, \omega, 1+\omega+u\omega, 1+u+\omega\right)$$

are respectively TypeI and Type II codes over $\mathbb{F}_4 + u\mathbb{F}_4$.

Proof. Self duality of the codes follows by Theorem 2.1 and the type of the code follows by the weights of the elements. For instance

$$\begin{aligned} & Q_p\left(u, 1+\omega, \omega+u\omega\right) Q_p\left(u, 1+\omega, \omega+u\omega\right)^T \\ = \; & Q_p(u^2 + (1+\omega)^2 + (\omega+u\omega)^2, u(1+\omega) + u(\omega+u\omega) + (1+\omega)(\omega+u\omega), \\ & u(1+\omega) + u(\omega+u\omega) + (1+\omega)(\omega+u\omega)) \\ = \; & Q_p(1,0,0) = I_p. \end{aligned}$$

Hence, the code $\mathcal{P}_p\left(u, 1+\omega, \omega+u\omega\right)$ is self-dual. In addition, $wt(1) = 2$, $wt(u) = 4$, $wt(1+\omega) = 1$ and $wt(\omega+u\omega) = 2$ so the weight of any row of $P_p\left(u, 1+\omega, \omega+u\omega\right)$ is $2 + 4 + \frac{p-1}{2}1 + \frac{p-1}{2}2 = 6 + 4k + 3 + 8k + 7 = 4(3k+4)$. Thus all codewords in

TABLE 1. Quadratic double circulant codes over $\mathbb{F}_4 + u\mathbb{F}_4$

the code over $\mathbb{F}_4 + u\mathbb{F}_4$	the binary image; $\phi \circ \psi(C)$
$\mathcal{B}_3(1+\omega, \omega+u\omega, 1+\omega, \omega, 1+\omega+u\omega, 1+u+\omega)$	$[32,16,8]_2$ Type II
$\mathcal{B}_3(1, u+u\omega, 1, u+u\omega, 1+u+u\omega, 1+u+u\omega)$	$[32,16,8]_2$ Type I
$\mathcal{P}_7(u, 1+\omega, \omega+u\omega)$	$[56,28,12]_2$ Type II
$\mathcal{B}_7(1+u+u\omega, \omega+u\omega, 1+\omega, u, 1+u\omega, 1+u\omega)$	$[64,32,12]_2$ Type II
$\mathcal{B}_{11}(1+\omega, \omega+u\omega, 1+\omega, \omega, 1+\omega+u\omega, 1+u+\omega)$	$[96,48,12]_2$ Type II
$\mathcal{B}_{11}(1, u+u\omega, 1, u+u\omega, 1+u+u\omega, 1+u+u\omega)$	$[96,48,12]_2$ Type I

$\mathcal{P}_p(u, 1+\omega, \omega+u\omega)$ have weight divisible by 4. The proofs of the other cases are analogous and skipped. □

We finish this section by presenting some examples of quadratic double circulant codes over $\mathbb{F}_4 + u\mathbb{F}_4$ in Table 1.

3.2. **($\mathbb{F}_4 + u\mathbb{F}_4$)-lifts of quadratic double circulant code $\mathcal{P}_7(0, 1+\omega, \omega)$ over $\mathbb{F}_4$.** We consider the lifts of the QDC code $\mathcal{P}_7(0, 1+\omega, \omega)$ over $\mathbb{F}_4$ whose Gray image is a self-dual $[28,14,6]_2$ code. The code is lifted to $\mathbb{F}_4 + u\mathbb{F}_4$ and as a result double circulant codes with binary image $[56,28,10]_2$ are obtained.

In the following let $\mathcal{K}_i$ be the $\mathbb{F}_4 + u\mathbb{F}_4$-code generated by $[I_7|A_i]$ where A_i is the circulant matrix with first row R_i.

$\mathcal{K}_i$	R_i
$\mathcal{K}_1$	$(u\omega, 1+\omega+u\omega, 1+u+\omega, \omega+u\omega, 1+u+\omega+u\omega, \omega, u+\omega)$
$\mathcal{K}_2$	$(u+u\omega, 1+\omega+u\omega, 1+u+\omega+u\omega, u+\omega, 1+\omega, \omega, u+\omega+u\omega)$
$\mathcal{K}_3$	$(u\omega, 1+\omega, 1+u+\omega, \omega+u\omega, 1+\omega+u\omega, u+\omega+u\omega, \omega)$

The binary images are $[56,28,10]_2$ self-dual codes, these are extended in Section 4.1 in order to obtain new binary codes of length 58.

3.3. **($\mathbb{F}_4 + u\mathbb{F}_4$)-lifts of quadratic double circulant code $\mathcal{B}_7(1, \omega, 1+\omega, 0, 1, 1)$ over $\mathbb{F}_4$.** The possible weight enumerators of Type I extremal self-dual codes of length 64 are characterized in [3] as:

$$W_{64,1} = 1 + (1312 + 16\beta)y^{12} + (22016 - 64\beta)y^{14} + \cdots \text{ where } 14 \leq \beta \leq 104,$$
$$W_{64,2} = 1 + (1312 + 16\beta)y^{12} + (23040 - 64\beta)y^{14} + \cdots \text{.where } 0 \leq \beta \leq 277.$$

Recently, codes with β =29, 39, 53 and 60 in $W_{64,1}$ and codes with β =51, 58 in $W_{64,2}$ are constructed in [20] and a code with $\beta = 80$ in $W_{64,2}$ is constructed in [12]. Together with these the existence of such codes is now known for β =14, 18, 22, 25, 32, 36, 39, 44, 46, 53, 60, 64 in $W_{64,1}$ and for β =0, 1, 2, 4, 5, 6, 8, 9, 10, 12, 13, 14, 16, 17, 18, 20, 21, 22, 23, 24, 25, 28, 29, 30, 32, 33, 36, 37, 38, 40, 41, 44, 48, 51, 52, 56, 58, 64, 72, 80, 88, 96, 104, 108, 112, 114, 118, 120, 184 in $W_{64,2}$.

The code $\mathcal{B}_7(1, \omega, 1+\omega, 0, 1, 1)$ over $\mathbb{F}_4$ has a binary image as the $[32,16,8]_2$ self-dual code with an automorphism group of order $2^{15} \times 3^2 \times 5 \times 7$. The code is lifted to $\mathbb{F}_4 + u\mathbb{F}_4$ and as binary image of the lifts extremal Type I codes of length 64 listed in Table 2 are obtained. In Table 2 $\mathcal{L}_i$ is the code over $\mathbb{F}_4 + u\mathbb{F}_4$ generated by $G_i = \left[\begin{array}{c|cc} I_8 & a & b \times \mathbf{1} \\ & c \times \mathbf{1}^T & A_i \end{array}\right]$ where A_i is the circulant matrix with first row

TABLE 2. Codes in $W_{64,1}$ as lifts of $B_7(1, \omega, 1+\omega, 0, 1, 1)$ over $\mathbb{F}_4$

$\mathcal{L}$	R_i; first row of A_i	
	(a, b, c); the border	β
$\mathcal{L}_1$	$(1, u+\omega+u\omega, \omega, 1+\omega+u\omega, \omega+u\omega, 1+\omega, 1+u+\omega)$	
	$(u\omega, 1+u+u\omega, 1+u\omega)$	18
$\mathcal{L}_2$	$(1+u\omega, u+\omega, \omega, 1+u+\omega+u\omega, u+\omega+u\omega, 1+u+\omega, 1+\omega)$	
	$(u\omega, 1+u+u\omega, 1+u\omega)$	32
$\mathcal{L}_3$	$(1+u, \omega, \omega, 1+\omega, \omega, 1+u+\omega+u\omega, 1+u+\omega)$	
	$(u+u\omega, 1+u+u\omega, 1+u\omega)$	46
$\mathcal{L}_4$	$(1+u+u\omega, \omega+u\omega, \omega, 1+\omega+u\omega, u+\omega, 1+\omega, 1+u+\omega)$	
	$(u+u\omega, 1+u\omega, 1+u\omega)$	60

R_i and $\mathbf{1}$ is the all 1 row vector of length 7 and the binary code $\phi \circ \psi(\mathcal{L}_i)$ has weight enumerator with β_i in $W_{64,1}$.

Remark 3.2. The code with $\beta = 60$ in $W_{64,1}$ was recently constructed in [20] and all the codes in Table 2 have an automorphism group of order $2^2 7$.

4. New extremal binary self-dual codes of lengths 58, 66 and 68

In this section, $\mathbb{F}_2$ and $(\mathbb{F}_2 + u\mathbb{F}_2)$-extensions of the codes constructed in the previous section are investigated. We were able to obtain 7 new binary codes of length 58, 24 new binary codes of length 66 and 29 new binary codes of length 68. All these codes have new weight enumerators.

4.1. New extremal binary self-dual codes of lengths 58.

The possible weight enumerators of an extremal self-dual binary code of length 58 are characterized in [3] as follows:

$$W_{58,1} = 1 + (165 - 2\beta)y^{10} + (5078 + 2\beta)y^{12} + \cdots \text{ where } 0 \le \beta \le 82$$
$$W_{58,2} = 1 + (319 - 24\beta - 2\gamma)y^{10} + \cdots \text{ where } 0 \le \beta \le 11 \text{ and } 0 \le \gamma \le 159 - 2\beta.$$

Recently, Yankov and Lee have presented the known extremal binary self-dual codes of length 58 and they obtain new ones in [19]. By considering lifts of $[8, 4, 4]_2$ Hamming code Karadeniz and Kaya constructed 11 new codes in $W_{58,2}$ in [9]. Together with the ones added from [19] and [9], the existence of such codes is known for $\beta = 55$ in $W_{58,1}$ and for $\beta = 0$ with $\gamma \in \{2m | m =$0, 1, 8, 9, 10, 15, 16, 34, 71, 79 or $18 \le m \le 64\}$, $\beta = 1$ with $\gamma \in \{2m | 21 \le m \le 57\}$ and $\beta = 2$ with $\gamma \in \{2m | m =$16, 18, 20, 21, 22, 19, 46, 49 or $24 \le m \le 44\}$ in $W_{58,2}$.

In this section, we obtain the codes in $W_{58,2}$ for $\beta = 0$ with $\gamma = 28$, $\beta = 1$ with $\gamma = 28, 32, 34, 38, 40$ and for $\beta = 2$ with $\gamma = 46$.

The binary Gray images of the codes $\mathcal{K}_1, \mathcal{K}_2$ and $\mathcal{K}_3$ are self-dual $[56, 28, 10]_2$ codes and when they are extended by Theorem 2.5 seven extremal self-dual codes of length 58 with new weight enumerators are obtained which are listed in Table 3.

TABLE 3. New codes in $W_{58,2}$ by Theorem 2.5 (7 codes)

$\mathcal{K}$	$X = (x_1, x_2, ..., x_{56})$	β	γ
$\mathcal{K}_1$	10011100000111010110110110010100100100010011110101110000	1	32
$\mathcal{K}_1$	10010010011100001101111010001111110011001111000101001111	1	38
$\mathcal{K}_1$	00111010100110000101110000110100011111011000001011001000	1	40
$\mathcal{K}_1$	00111111011100001001000010010000010000010000101010001000	2	46
$\mathcal{K}_2$	11010000110101000111011001001100010010100011100111101010	0	28
$\mathcal{K}_2$	01101011110110000100001101010100111100100100110110010111	1	34
$\mathcal{K}_3$	01110000010011000101111110000111010100000010101001100110	1	28

TABLE 4. New codes in $W_{66,3}$ by Theorem 2.6 (13 codes)

C	$X = (x_1, x_2, ..., x_{32})$	β
$\mathcal{L}_1$	01101010101010101101101110100100	35
$\mathcal{L}_1$	00100101110100100000100110110001	36
$\mathcal{L}_1$	10110000001001110101100111100001	37
$\mathcal{L}_1$	11100001011100101111010110010001	38
$\mathcal{L}_2$	01011001111111001001101010111001	43
$\mathcal{L}_2$	10110110001011100111101100111001	44
$\mathcal{L}_2$	00000010110011001100010111000011	47
$\mathcal{L}_2$	01101001101001111110011111100100	48
$\mathcal{L}_2$	10010010110101101101100111011110	51
$\mathcal{L}_3$	11001001111101100011101001000101	60
$\mathcal{L}_3$	01101001010111101100000011110011	67
$\mathcal{L}_4$	01010111010000001100100111001110	70
$\mathcal{L}_4$	00101101101000101100010011101100	75

4.2. **New extremal binary self dual codes of length** 66. A self-dual $[66, 33, 12]_2$-code has a weight enumerator in one of the following forms ([4])

$$\begin{aligned} W_{66,1} &= 1 + (858 + 8\beta)\, y^{12} + (18678 - 24\beta)\, y^{14} + \cdots \text{ where } 0 \le \beta \le 778, \\ W_{66,2} &= 1 + 1690y^{12} + 7990y^{14} + \cdots \\ \text{and } W_{66,3} &= 1 + (858 + 8\beta)\, y^{12} + (18166 - 24\beta)\, y^{14} + \cdots \text{ where } 14 \le \beta \le 756, \end{aligned}$$

Recently, five new codes in $W_{66,1}$ are constructed in [12]. For a list of known codes in $W_{66,1}$ we refer to [12, 18] and references therein.

First codes with a weight enumerator in $W_{66,3}$ were discovered in [18] and recently 14 codes were discovered in [10] together with these the existence of codes in $W_{66,3}$ is known for β =28, 29, 30, 31, 32, 33, 34, 49, 50, 54, 55, 56, 57, 58, 59, 62, 63 and 66.

In this work, we construct 24 codes with new weight enumerators, more precisely the codes with β =35, 36, 37, 38, 43, 44, 47, 48, 51, 60, 67, 70, 71, 73, 74, 75, 76, 77, 78, 79 and 80 in $W_{66,3}$.

Remark 4.1. In order to apply the extension in Theorem 2.6 the generator matrices for the binary image of $\mathcal{L}_i$ are converted into standard form $[I_{32}|A_i]$ and the matrices are available online at [13]. So, in Table 4 the codes are generated by the matrices

TABLE 5. New codes in $W_{66,3}$ by Theorem 2.5 (11 codes)

C	$X = (x_1, x_2, ..., x_{64})$	β
$\mathcal{L}_2$	1100101011011100011110010011111101101001011100101110010100111111	45
$\mathcal{L}_2$	0011110011001011010001000001101110011100101111000010000111011100	46
$\mathcal{L}_3$	0000100000001000110000011100010000001011101000001000001011110011	61
$\mathcal{L}_4$	0001101001010010010010111101111101111110010111100001000011010001	71
$\mathcal{L}_4$	1111101001110110001000111011111101101101101101100001111101101110	73
$\mathcal{L}_4$	1110011001010110000001010111001000001110011111010101110110010101	74
$\mathcal{L}_4$	1010110101111010100111100111010001100000011100101110000000011001	76
$\mathcal{L}_4$	0111111111011001001011011001001111010101011001011110110101000010	77
$\mathcal{L}_4$	1100101100011101111011110101010011000010010101011110001111011111	78
$\mathcal{L}_4$	1011110101001100001100101110011000001110100011010110001011110011	79
$\mathcal{L}_4$	0000100100100100111100111011011010100100000001000101100111110110	80

of the form

$$\left[\begin{array}{cc|ccccccc} 1 & 0 & x_1 & \dots & x_{32} & 1 & \dots & 1 \\ \hline y_1 & y_1 & & & & & & \\ \vdots & \vdots & & & I_{32} & & & A_i \\ y_{32} & y_{32} & & & & & & \end{array}\right].$$

In addition, as binary extensions by Theorem 2.5 we obtained 11 new codes in $W_{66,3}$ which are listed in Table 5.

Remark 4.2. In Tables 3 and 5 the codes are generated by the matrices of the form

$$\left[\begin{array}{cc|c} 1 & 0 & X \\ \hline y_1 & y_1 & \\ \vdots & \vdots & \phi \circ \psi\,(C) \\ y_k & y_k & \end{array}\right].$$

4.3. **New extremal binary self dual codes of length** 68**.** The weight enumerator of an extremal binary self-dual code code of length 68 is in one of the following forms ([4]):

$$\begin{aligned} W_{68,1} &= 1 + (442 + 4\beta)\, y^{12} + (10864 - 8\beta)\, y^{14} + \cdots \text{ where } 104 \le \beta \le 1358, \\ W_{68,2} &= 1 + (442 + 4\beta)\, y^{12} + (14960 - 8\beta - 256\gamma)\, y^{14} + \cdots \end{aligned}$$

where $0 \le \gamma \le 11$ and $14\gamma \le \beta \le 1870 - 32\gamma$. Tsai et al. constructed a substantial number of codes in both possible weight enumerators in [18]. Recently, 178 new codes are obtained in [15] and 28 new codes including the first examples with $\gamma = 4$ and $\gamma = 6$ in $W_{68,2}$ are obtained in [11]. Together with the ones in [11, 15] codes exists for $W_{68,2}$ when $\gamma = 0$ and β =38, 40, 44,...,139, 141, 142, 143, 145, 147, 148, 149, 151, 153, 170, 204, 238, 272; $\gamma = 1$ and β =54, 56, 58, 60, ..., 66, 68, 70, 72,..., 115, 118, 119, 120, 123, 125, 126, 129, 132, 133, 135, 137, ...,149, 150, 151, 153, 155, 159; $\gamma = 2$ and β =65, 68, 71, 77, 85, 87, 89, 91, 93, 95, 97, 99, 101, 103, 105, 109, 111, 115, 117, 119, 121, 123, 125, 127, 129, 131, 133, 135, 137, 139, 145, 151, 153, 155, 158, 160, 162 or $\beta \in \{2m | 37 \le m \le 68,\ 70 \le m \le 76\}$; $\gamma = 3$ and β =88, 90, 96, 100, 102, 104, 108, 112, 114, 116, 117, 126, 127, 128, 130, 133, 136, 137, 138,

TABLE 6. New codes in $W_{68,2}$ by Theorem 2.5 for $\mathbb{F}_2 + u\mathbb{F}_2$ (29 codes)

C	$X = (x_1, x_2, ..., x_{32})$	c	γ	β
$\mathcal{L}_1$	$[0300u01333u330uuu10011100131u111]$	$1+u$	1	67
$\mathcal{L}_1$	$[33010u0u003u11u013331uu13u130031]$	1	1	69
$\mathcal{L}_1$	$[u311uu0uu0101u01u0000u11310uu1u0]$	1	1	71
$\mathcal{L}_1$	$[3u0uu1u33030110031u1330u0011uuu1]$	$1+u$	2	107
$\mathcal{L}_1$	$[013311u1uuu1u103uu111300u3100u31]$	1	3	101
$\mathcal{L}_1$	$[00311u000u0uu101u301u13030111000]$	1	3	110
$\mathcal{L}_1$	$[0u111u000u0003u103u3u11030131u0u]$	1	3	122
$\mathcal{L}_1$	$[111u010u0u003u113001010103111031]$	1	3	123
$\mathcal{L}_2$	$[u11u0303u3u0113330uuu3031110u130]$	$1+u$	1	117
$\mathcal{L}_2$	$[30033u31u30301133uu13u311311101u]$	1	2	113
$\mathcal{L}_2$	$[11101u3001100uu3u133u1u3001u1130]$	$1+u$	3	132
$\mathcal{L}_3$	$[uu3113000031u11310011333u3u13u10]$	1	1	116
$\mathcal{L}_3$	$[30113uuuuu33uu03u1u3uu311u00u0u1]$	1	1	124
$\mathcal{L}_3$	$[313310uu3010u3003u003030u3031003]$	$1+u$	1	134
$\mathcal{L}_4$	$[1003u10u1030u10101u1001u00uuuu1u]$	1	1	121
$\mathcal{L}_4$	$[1000133u3010013uu13uuu11u031uuu1]$	1	1	122
$\mathcal{L}_4$	$[130u31133u31uu30u100u03u3110u31u]$	$1+u$	1	127
$\mathcal{L}_4$	$[101103310u133003000033u3uu111uu1]$	$1+u$	1	128
$\mathcal{L}_4$	$[30u103313u031100u13uu101u03011u3]$	1	1	130
$\mathcal{L}_4$	$[1u1u10uu0uu0u31u0u103u133u11333u]$	$1+u$	1	131
$\mathcal{L}_4$	$[10u011u001u310300330u03uu3uuu101]$	$1+u$	1	136
$\mathcal{L}_4$	$[330u0113u0uu3333u003u000uu010u33]$	1	1	157
$\mathcal{L}_4$	$[011uu103103111u00030031u0110331u]$	1	2	143
$\mathcal{L}_4$	$[00u1u0u1u1u30130u31u030011u00130]$	$1+u$	2	147
$\mathcal{L}_4$	$[uuu10uu30103u310u310u3uu1100u31u]$	$1+u$	2	149
$\mathcal{L}_4$	$[33uuu0u3uu110330u3010u110301u101]$	$1+u$	2	154
$\mathcal{L}_4$	$[330uu0030033031001010u310103u1u1]$	1	2	156
$\mathcal{L}_4$	$[0001uuu3u3u3u3100330u3001100033u]$	1	2	159
$\mathcal{L}_4$	$[3031u111133u01u133u1u00113303301]$	$1+u$	3	156

140, 141, 142, 144, 145, 147, 148, 149, 153, 154, 158, 159, 160, 162, 176, 188, 193, 196; $\gamma = 4$ and β =102, 110, 116, 120, 122, 124, 128, 130, 134, 136, 138, 140, 142, 150, 152, 154, 156, 158, 160, 162, 164, 166, 168, 170, 172, 174, 176, 180 and $\gamma = 6$ with β =138, 154, 156, 158, 162, 176, . For a list of known codes in $W_{68,1}$ we refer to [18].

In this work as binary images of $(\mathbb{F}_2 + u\mathbb{F}_2)$-extensions of codes in Table 6 we obtain 29 codes with new weight enumerators in $W_{68,2}$, more precisely codes with $\gamma = 1$, β =67, 69, 71, 116, 117, 121, 122, 124, 127, 128, 130, 131, 134, 136, 157; with $\gamma = 2$, β =107, 113, 143, 147, 149, 154, 156, 159 and codes with rare parameters $\gamma = 3$, β =101, 110, 122, 123, 132, 156.

Remark 4.3. In Table 6 for the extension vectors X over $\mathbb{F}_2 + u\mathbb{F}_2$ the element $1+u$ is abbreviated as 3 and the codes are generated by the matrices of the following

form

$$G = \left[\begin{array}{cc|c} 1 & 0 & X \\ \hline y_1 & cy_1 & \\ \vdots & \vdots & \psi(\mathcal{L}_i) \\ y_k & cy_k & \end{array}\right].$$

The binary code generated by $\phi(G)$ has a new weight enumerator in $W_{68,2}$ for the given parameters. The binary generator matrices for codes with $\gamma = 3$ in $W_{68,2}$ in Table 6 are available online at [13].

References

[1] K. Betsumiya, S. Ling and F. R. Nemenzo, "Type II codes over $\mathbb{F}_{2^m} + u\mathbb{F}_{2^m}$", *Discrete Mathematics*, Vol. 275 pp. 43-65, 2004.

[2] W. Bosma, J. Cannon, and C. Playoust, The Magma algebra system. I. The user language, *J. Symbolic Comput.*, Vol 24, pp. 235–265, 1997.

[3] J. H. Conway, and N. J. A. Sloane, "A new upper bound on the minimal distance of self-dual codes", *IEEE Trans. Inform. Theory*, Vol. 36, 6, 1319–1333, 1990.

[4] S. T. Dougherty, T. A. Gulliver, and M. Harada,"Extremal binary self dual codes", *IEEE Trans. Inform. Theory*, Vol. 43 pp. 2036-2047, 1997.

[5] S. T. Dougherty, J-L. Kim and P, Sole, "Double circulant codes from two class association schemes", *Advances in Mathematics of Communications*, Vol. 1, No.1, pp. 45-64, 2007.

[6] P. Gaborit, "Quadratic double circulant codes over fields", *Journal of combinatorial theory*, Vol. 97, Issue 1, pp. 85-107, 2002.

[7] T. A. Gulliver and M. Harada, "On double circulant doubly even self-dual $[72, 36, 12]$ codes and their neighbors ", *Australasian Journal of Combinatorics*, Vol. 40, 137–144, 2008.

[8] W.C. Huffman and V. Pless, "Fundamentals of error correcting codes", *Cambridge University press*, 2003.

[9] S. Karadeniz and A. Kaya, "New extremal binary self-dual codes of length 58 as R_3-lifts from the shortened binary $[8, 4, 4]$ Hamming code", *J. Franklin Inst.,* Vol. 349, No. 9, pp. 2824–2833, 2012.

[10] S. Karadeniz and B. Yildiz, "New extremal binary self-dual codes of length 66 as extensions of self-dual codes over R_k", *J. Franklin Inst.,* Vol. 350, No. 8, pp. 1963–1973, 2013.

[11] S. Karadeniz and B. Yildiz, "New extremal binary self-dual codes of length 68 from R_2-lifts of binary self-dual codes ", *Advances in Mathematics of Communications*, Vol. 7, No. 2, pp. 219–229, 2013.

[12] S. Karadeniz, B. Yıldız and N., Aydın, "Extremal binary self-dual codes of lengths 64 and 66 from four-circulant constructions over codes $\mathbb{F}_2 + u\mathbb{F}_2$", *to appear in FILOMAT.*

[13] A. Kaya, B. Yildiz, and İ. Şiap, *Generator matrices for new extremal self-dual binary codes*, available online at http://www.fatih.edu.tr/~akaya/QDC.html

[14] A. Kaya, B. Yildiz and I. Siap, "New extremal binary self-dual codes of length 68 from quadratic residue codes over $\mathbb{F}_2 + u\mathbb{F}_2 + u^2\mathbb{F}_2$", **to appear in** Finite Fields Appl, 2014.

[15] A. Kaya and B., Yildiz, "Extension theorems for self-dual codes over rings and new binary self-dual codes", available online at http://arxiv.org/abs/1404.0195.

[16] S. Ling and P. Sole, "Type II codes over $\mathbb{F}_4 + u\mathbb{F}_4$", *Europ. J. Combinatorics*, Vol. 22 pp. 983-997, 2001.

[17] E. M. Rains, "Shadow Bounds for Self Dual Codes", *IEEE Trans. Inf. Theory*, Vol.44, pp.134–139, 1998.

[18] H-P. Tsai, P-Y. Shih, R-Y. Wuh, W-K. Su and C-H. Chen, "Construction of self-dual codes", *IEEE Trans. Inform. Theory*, Vol. 54, pp. 3826–3831, 2008.

[19] N. Yankov, M-H. Lee, "New binary self-dual codes of lengths 50-60", **to appear in** *Des. Codes Cryptogr*, 2013.

[20] N. Yankov, "Self-dual $[62, 31, 12]$ and $[64, 32, 12]$ codes with an automorphism of order 7", *Advances in Mathematics of Communications*, Vol.8, No1. pp.73–81, 2014.

Department of Mathematics, Fatih University, 34500, Istanbul, Turkey
E-mail address: byildiz@fatih.edu.tr, akaya@fatih.edu.tr

Department of Mathematics, Yildiz Technical University, 34210, Istanbul, Turkey
E-mail address: isiap@yildiz.edu.tr